

HANDBOOK ON

REFERENCE

METHODS

SOIL

TESTING



**THE COUNCIL ON
SOIL TESTING AND
PLANT ANALYSIS**



Table 2. SOIL TEST PHOSPHORUS METHODS


Parameter	Method				
	Mehlich No. 1 (Double Acid)	Bray P1	Mehlich No. 2	Olsen	AB-DTPA
Adaptability Limits	Sandy soils, acid, low in CEC	Acid soils with moderate CEC	All soils	Alkaline soils	Alkaline soils
Sample Size, Weight (volume)	5 g (4-cm ³)	2 g (1.70-cm ³)	2.5-cm ³	2.5 g (2-cm ³)	10 g (8.5-cm ³)
Volume of Extractant, ml	25	20	25	50	20
Extracting Solution	0.05N HCl in 0.025N H ₂ SO ₄	0.03N NH ₄ F in 0.025N HCl	0.2N HOAc in 0.015N NH ₄ F in 0.2N NH ₄ Cl in 0.012N HCl	0.5N NaHCO ₃ at pH 8.5	1M NH ₄ HCO ₃ 0.005M DTPA at pH 7.6
Shaking Time, minutes	5	5	5	30	15
Shaking Action and Speed	Reciprocating 180+ oscillations/min.	Reciprocating 180+ oscillations/min.	Reciprocating 180+ oscillations/min.	Reciprocating 180+ oscillations/min.	Reciprocating 180+ oscillations/min.
Method of P Determination in Extract	Molybdenum Blue	Molybdenum Blue	Molybdenum Blue	Molybdenum Blue	Molybdenum Blue
Range in Soil P concn without Dilution, kg/ha	2-100	2-250	2-200	2-200	2-100
Sensitivity, kg/ha	1	1	1	1	2
Primary Reference	North Carolina Soil Test Div. Mimeo (1953)	Soil Sci. 59:39 (1945)	Comm. Soil Sci. Plant Anal. 9 (6):477 (1978)	USDA Circular 939 (1954)	Comm. Soil Sci. Plant Anal 8:195 (1977)

Table 3. SOIL TEST POTASSIUM METHODS

340
159

Circular No. 939

March 1954 • Washington, D. C.



UNITED STATES DEPARTMENT OF AGRICULTURE

Estimation of Available Phosphorus in Soils by Extraction With Sodium Bicarbonate¹

By STERLING R. OLSEN, *senior soil scientist*, C. V. COLE, *soil scientist*, FRANK S. WATANABE, *soil scientist*, and L. A. DEAN, *principal soil scientist*, Soil and Water Conservation Research Branch, Agricultural Research Service²

United States Department of Agriculture in cooperation with the Colorado Agricultural Experiment Station

CONTENTS

	Page		Page
Introduction.....	1	Evaluation of method—Con.	
Development of method.....	2	Relation of phosphate-fertility	
Theoretical basis.....	2	level to yield response.....	9
Extraction details.....	4	Greenhouse experiments....	9
Time of extraction.....	4	Field experiments.....	11
pH of extractant.....	5	Measurement of NaHCO ₃ -	
Method of extraction.....	5	soluble phosphorus in rela-	
Evaluation of method.....	7	tion to yield response.....	12
Soil samples.....	7	Correlation of 'A' values with	
Experiment 1.....	7	various methods.....	13
Experiment 2.....	8	Discussion.....	15
Experiment 3.....	8	Recommendations.....	17
Experiment 4.....	8	Summary.....	17
Experiment 5.....	8	Literature cited.....	18
Experiment 6.....	9		

INTRODUCTION

Relatively few methods employing either an alkaline or acid extractant have been proposed for measuring readily available phosphorus on both acid and alkaline soils. Das (9)³ recommended a 1-percent K₂CO₃ solution for extracting phosphates on both acid and calcareous soils. Hockensmith and coworkers (14) modified this method and reported that K₂CO₃ is the most satisfactory of several extractants studied for indicating the crop response to applied phosphate on calcareous soils. On acid soils Gardner and Kelley (13) reported that

¹ Approved by the director of the Colorado Agricultural Experiment Station as Scientific Journal Series No. 418.

² The authors wish to express appreciation for the many helpful suggestions given by Robert Gardner, professor of agronomy, Colorado Agricultural Experiment Station, during the course of these studies. Credit is extended also to various staff members in other State stations who conducted many of the field experiments and made available the samples for the greenhouse studies.

³ Italic numbers in parentheses refer to Literature Cited, p. 18.

the K_2CO_3 method was unsuitable, since it extracted relatively much more phosphorus than was available to plants.

Rhoades (23) leached soils with a solution of 0.01 N Na_2CO_3 buffered with sufficient H_3BO_3 added to give it the same pH as the soil. For most calcareous soils this solution was largely a mixture of sodium bicarbonate and sodium borate with pH values between 7 and 8.5. This method was not proposed for acid soils.

A new method for extracting soil phosphorus with sodium bicarbonate solutions is described in this circular. The relationship of the extractable phosphorus to yield response in greenhouse and field experiments and to a measure of plant available phosphorus, or the 'A' value (12), is shown. The $NaHCO_3$ -soluble phosphorus is compared with the methods of Bray (4), Truog, CO_2 -soluble (26), and H_2O -soluble (2) by means of correlation analysis with 'A' values.

DEVELOPMENT OF METHOD

THEORETICAL BASIS

Several studies have been conducted to show the relationship of pH to phosphate solubility in calcareous soils (6, 13, 22, 26) and in systems containing excess $CaCO_3$ (1, 7). The results show that the solubility of the calcium phosphate is a function of pH and Ca^{++} activity, with a minimum solubility between pH 7 to 7.5 and an increase in solubility on both the alkaline and acid side of this pH range. On the alkaline side the solubility of phosphorus in the presence of solid phase $CaCO_3$ is a function of the Ca^{++} activity, whereas on the acid side of the minimum solubility range in pH the solubility of phosphorus is a function of both the H^+ activity and the Ca^{++} activity (7). An increase in H^+ activity increases the solubility of phosphorus and an increase in Ca^{++} activity decreases the solubility of phosphorus.

One of the objectives in research on soil tests has been to eliminate or diminish the secondary precipitation reactions that may occur during the extraction process with dilute acids (8). McGeorge and others (18) stated that the solvent action of CO_2 on phosphorus in calcareous soils was mainly a function of reaction and that the H^+ activity must be sufficient to overcome the common ion effect of Ca^{++} in precipitating a calcium phosphate. Thus, two reactions with opposite effects on phosphate solubility occur when CO_2 is bubbled through a calcareous soil:water suspension, i. e., an increase in solubility caused by a decrease in pH and a decrease in solubility when precipitation of a calcium phosphate occurs as a result of the increase in Ca^{++} activity. Stanberry (26) showed that both reactions affected phosphate solubility on a calcareous Sagemoor fine sandy loam, since the concentration of phosphorus was higher at both pH 8.5 and 5.8 than at intermediate pH values. Time-of-extraction studies on this same soil with CO_2 showed that there was a maximum solubility of phosphorus in 5 and 30 minutes, respectively, for low and high phosphate fertility levels. A decrease in solubility of phosphate following the maximum level led to the conclusion that a calcium phosphate precipitated from supersaturated solutions of the CO_2 extract.

Secondary precipitation reactions may occur with extractants that increase the H^+ activity of the calcareous soil:water system, because a concomitant increase in the Ca^{++} activity occurs. Since the solubility of phosphate may be increased in this system with a decrease in Ca^{++} activity, the secondary precipitation reactions should be largely eliminated with an extractant that decreases the Ca^{++} activity.

Gardner and Kelley (13) indicated that in the presence of $CaCO_3$ and excess carbonate ions the increase in phosphorus solubility in 1-percent K_2CO_3 solutions on calcareous soils was a result of the repression of the Ca^{++} activity. The K_2CO_3 method is not applicable to acid soils, however, probably because forms of iron and aluminum phosphates unavailable to plants are dissolved as a result of the high pH of the extracting solution.

The Ca^{++} activity in a soil:water system is repressed with $NaHCO_3$ solutions at pH 8.5, and the undesirable effects of the high pH of 1-percent K_2CO_3 solutions are avoided. The ratio of HCO_3^- to CO_3^{--} is 63 to 1 in a solution of $NaHCO_3$ at pH 8.5. The influence of $NaHCO_3$ concentration on the molar concentrations of HCO_3^- , CO_3^{--} , and Ca^{++} in a $CaCO_3-H_2O$ system at pH 8.5 is shown in table 1. The calculations are based on the mass-action law, using 4.82×10^{-9} as the solubility product of calcite (11) with no corrections for salt effect. In a 0.5 molar solution of $NaHCO_3$, the equilibrium calcium concentration is 857 times smaller than in a pure $CaCO_3-H_2O$ system in equilibrium with atmospheric CO_2 . The main effect of the $NaHCO_3$ in calcareous soils, therefore, is to decrease the Ca^{++} activity, which in turn increases the solubility of phosphorus.

In acid and neutral soils the phosphorus present in the form of calcium phosphates would probably become more soluble in the presence of $NaHCO_3$, owing to a repression of the Ca^{++} activity, assuming that solid phase $CaCO_3$ would be formed at pH 8.5. Since the calcium phosphates are known to be the most available forms of phosphorus in these soils, the $NaHCO_3$ method may be applicable to acid and neutral as well as to calcareous soils.

The main effect of the $NaHCO_3$ on acid and neutral soils probably would be through ionic competition of HCO_3^- , CO_3^{--} , and OH^- ions for phosphate adsorbed on the surface of soil particles. According to Kurtz and others (16) the replacing ability of HCO_3^- for phosphate was greater than the replacing ability of acetate or sulfate ions on acid soils. Laatsch (17) stated that HCO_3^- ions displaced sorbed phosphate from colloidal surfaces in soils.

TABLE 1.—Influence of $NaHCO_3$ concentration on the molar concentrations of HCO_3^- , CO_3^{--} , and Ca^{++} in a $CaCO_3-H_2O$ system at pH 8.5

$NaHCO_3$	HCO_3^-	CO_3^{--}	Ca^{++}
¹ 0.0	0.0014	2.30×10^{-5}	5.30×10^{-4}
.01	.0098	1.56×10^{-4}	3.09×10^{-5}
.10	.0984	1.56×10^{-3}	3.09×10^{-6}
.50	.492	7.80×10^{-3}	6.18×10^{-7}

¹ Solubility of calcite in H_2O at atmospheric pressure of CO_2 (11).

The amount of phosphorus on the surface of soil particles, which readily exchanges with P^{32} in the soil solution, was shown to be highly correlated with plant available phosphorus, or 'A' values (21).

The relationship between the surface or exchangeable phosphorus measured by P^{32} and the phosphorus soluble in $NaHCO_3$ is shown in




FIGURE 1.—Relationship between surface phosphorus and phosphorus soluble in 0.5 molar $NaHCO_3$ solutions on 25 soils.

figure 1 for 25 soils. The pH of these soils ranged from 5.0 to 7.6. The $NaHCO_3$ extracts about 50 percent of the surface phosphorus. The data indicate that the $NaHCO_3$ -soluble phosphorus is closely related to a form of phosphorus, which in turn is highly correlated with the amount of phosphorus available to plants.

EXTRACTION DETAILS

Time of Extraction

Five soil types, representing acid to highly calcareous soils, were selected for this study (table 2). Samples from plots receiving applications of phosphate in the field are included to illustrate the effect of phosphate-fertility level. The extracting solution was 0.5 molar $NaHCO_3$ solution at pH 8.5 and the soil:solution ratio was 1:20.

Equilibrium is not reached in any of the samples over a 2-hour extraction period (fig. 2); however, the curves tend to level off after 30 minutes on the low-phosphate-fertility soils. The absolute differences between the low- and high-phosphate-fertility levels become greater as the extraction time increases; the percentage difference, however, remains about the same for all extraction periods studied. The correlation coefficients between 'A' values and NaHCO_3 -soluble phosphorus were essentially the same for 30-, 60-, and 120-minute extraction periods for experiments 4 and 5 described later. Correlations for shorter extraction periods were not determined.

TABLE 2.—*Calcium carbonate content, pH, and phosphate treatment of soil types used for time-of-extraction studies*

Soil No.	Soil type and treatment	CaCO_3	pH
		<i>Percent</i>	
326	Davidson silty clay loam ¹		5.3
318	Cecil loam ¹		5.7
333	Owyhee silt loam		7.0
335	Owyhee silt loam, plus phosphate		7.1
151	Pryor silty clay	0.51	7.7
145	Pryor silty clay, plus manure	1.15	7.6
428	Millville loam	55.8	8.2
433	Millville loam, plus phosphate	55.8	8.2

¹ Data on yield response of soils to phosphate in the greenhouse indicated that the Cecil loam had a higher phosphate-fertility level than the Davidson silty clay loam.

pH of Extractant

An increase in the pH of the extracting solution from 8.5 to 9.0 causes an increase in phosphorus solubility. The effect of the pH increase is greater on acid than on alkaline soils (table 3). Correlation coefficients for the phosphate soluble in NaHCO_3 compared with 'A' values showed very little difference between pH 8.5 and 9.0. The relative differences in the extractable phosphorus are less at pH 8.5 than at pH 9 as the soil pH changes (table 3). The level of extractable phosphorus in relation to crop response, therefore, would be more uniform at pH 8.5 than at pH 9 for a group of soils varying in pH.

Method of Extraction

The following method is recommended for extracting soils with NaHCO_3 solutions: (a) Prepare a 0.5 molar NaHCO_3 solution adjusted to pH 8.5 with NaOH . (b) Shake 5 gm. of soil with 100 ml. of the NaHCO_3 solution and 1 teaspoon of carbon black (Darco G-60 or other suitable carbon black) for 30 minutes. The rate of shaking should be constant. Add more carbon black to the flask after shaking if filtrate is not clear. (c) Filter through Whatman No. 40 or other suitable paper. (d) Add 5 ml. of the filtrate to a 25-ml. volumetric flask and determine phosphorus by the Dickman and Bray method (10). The molybdate reagent is modified by adding an extra 50 ml. of concentrated HCl per liter to neutralize the NaHCO_3 in a 5-ml.




FIGURE 2.—Relation of effect of time of extraction on NaHCO_3 -soluble phosphorus in pounds P_2O_5 per acre from several soil types to phosphate-fertility level.

aliquot. (e) On some soils high in extractable phosphorus it may be necessary to take less than a 5-ml. aliquot. If so, the total volume should be made up equivalent to 5 ml. of the NaHCO_3 solution in order to maintain the proper acidity for developing the color. The standard curve should be prepared with the NaHCO_3 included. (f) The carbon black (Darco G-60) releases phosphorus when treated with NaHCO_3 . Therefore, the carbon black is pretreated once with the NaHCO_3 solution by leaching, washing with water, and drying.

TABLE 3.—Effect of pH of the NaHCO_3 extracting solution on the mean amount of phosphate extracted from soils varying in pH

Number of soils	pH range of soils	Pounds P_2O_5 per acre soluble in NaHCO_3 at—	
		pH 8.5	pH 9.0
41-----	4.5 to 6.0	101	150
23-----	6.1 to 7.0	71	89
10-----	7.1 to 7.6	46	52

It may be well to indicate a few precautions to be observed in extracting soils. If the flask containing the aliquot is shaken too vigorously after the addition of the molybdate reagent, some solution may be lost as a result of rapid evolution of CO_2 . After CO_2 evolution has subsided, wash down the neck of the flask and dilute to about a 20-ml. volume. Add 1 ml. of the dilute SnCl_2 solution, shake immediately, dilute to volume, and mix well. Failure to observe this precaution with respect to the SnCl_2 will give erratic results. The pH of the NaHCO_3 solution tends to increase on standing if exposed to the atmosphere. Changes in pH of the NaHCO_3 solution can be prevented by the addition of mineral oil.

EVALUATION OF METHOD

SOIL SAMPLES

Laboratory and greenhouse studies were conducted on six groups of soil samples to obtain a measure of yield and 'A' value. Response data from field experiments were available from a part of these samples. The greenhouse studies on the six groups of soils are described by experiment number. The samples were collected from the surface 6 inches of soil. Nitrogen and potash were added in amounts considered to be sufficient for maximum growth in all experiments. The crops grown on these soils were harvested at the heading-out stage, except in experiment 6.

Experiment 1

Soil samples were collected from 74 locations in the United States and Saskatchewan, Canada. The pH values were less than 6 on 41 samples, between 6.1 and 7 on 23 samples, and above 7.1 on 10

samples. Free calcium carbonate was found in 12 samples. The experiment was conducted during the summer of 1951 at Beltsville, Md., to measure the yield and 'A' value for each soil.⁴ Superphosphate containing P³² was thoroughly mixed with the soil at the rate of 40, 80, and 160 pounds P₂O₅ per acre. The crop was German millet.

The 'A' values used for the correlation studies in both experiments 1 and 2 were the average values for the various rates of P₂O₅ added. With most soils the 'A' value was essentially the same for each rate of P₂O₅ added.

Experiment 2

An additional 40 soil samples were collected from various sections of the United States, and an experiment similar to the one described previously was conducted in the summer of 1952 at Beltsville, Md. The 80-pound P₂O₅ rate was omitted in this experiment. The pH values were less than 6 on 23 samples, between 6.1 and 7 on 5 samples, and above 7.1 on 12 samples. The crop was German millet.

Experiment 3

A group of 38 soils representing 28 different soil types was obtained from various locations in Colorado. Saturated paste pH values were between 6 and 7 for 3 soils and the rest ranged from pH 7 to 8.4. The calcium carbonate content ranged from none on the 3 acid soils to 9.6 percent. Phosphorus was mixed with the soil at the rate of 80 pounds of P₂O₅ per acre. The phosphorus in the form of H₂PO₄ containing P³² was adsorbed on a resin, IR-4B. A separate experiment on several soils showed that the phosphated resin acted essentially the same as superphosphate as a source of phosphorus for the plants. This experiment was conducted during the summer of 1952 at Fort Collins, Colo., using German millet as a crop.

Experiment 4

Thirty soil samples from long-term rotation plots at three locations were selected. One-half of the samples were from plots that had received either manure or phosphate, and the other half had received no phosphorus applications for 10 years or more. All these samples contained free calcium carbonate. The rate of application and the form of the phosphate were the same as in experiment 3. This experiment and experiment 5 were conducted at Fort Collins in the spring and fall of 1952, using oats as a crop.

Experiment 5

Thirty soil samples, representing 10 soil types and 3 levels of available phosphorus, were selected from 10 locations. The samples were obtained from field plots that had received 0, 120, and 240 pounds of P₂O₅ per acre 1 year previous to sampling time. All of the soil pH values were above 7, but 4 of the soil types did not contain any free calcium carbonate and the remaining 6 were calcareous.

⁴The greenhouse experiments 1 and 2 were conducted by Maurice Fried, Beltsville, Md.

Experiment 6

Thirty soil samples from as many different locations were collected from Oregon. Phosphorus was mixed with the soil at the rate of 100 pounds of P_2O_5 per acre. The phosphorus in the form of H_2PO_4 containing P^{32} was adsorbed on a resin, IRA-400. The crop was Sudan grass, and the experiment was conducted in the summer of 1952 at Corvallis, Oreg. The Sudan grass was harvested after 7 weeks' growth.⁵

Yield data from this experiment are not discussed, since only three soils showed a significant response to applied phosphate.

RELATION OF PHOSPHATE-FERTILITY LEVEL TO YIELD RESPONSE

The acceptable standard for determining the relative value of soil tests is the yield response of crops to applied phosphorus in greenhouse and field experiments. With the latter, especially, the assurance that the yield response measured was a result of differences in phosphate-fertility levels was not always evident. Furthermore, the relationship between phosphate-fertility level and the soluble phosphate by the soil test, on the soils showing no yield response of crops, was indeterminate.

The method of Fried and Dean (12) for determining the plant available phosphorus, or 'A' value, permits a measurement to be made both on soils responding and not responding to applied phosphorus; the relationship, therefore, between 'A' value and phosphate soluble in chemical extractants can be determined for a wide range of phosphate-fertility levels. The 'A' value is defined as the amount of soil phosphorus that is as available to the plants as the phosphorus in a standard phosphate fertilizer which has been mixed with the soil. Since the 'A' value is used in this study for comparing the relative value of various soil extractants for phosphorus, the relationship between 'A' value and yield response is shown.

Greenhouse Experiments

The relationship between the percentage of soils showing a significant (5-percent level) crop yield increase from applied phosphorus and the 'A' value is shown in figure 3. A total of 212 samples are compared.

Where 'A' values were below 50 pounds P_2O_5 per acre, all greenhouse soils showed crop yield increases from phosphate fertilization. In the range from 50 to 100 pounds P_2O_5 per acre, 98 percent of the soils showed a crop response to phosphorus. As the 'A' values increased the proportion of soils showing response decreased rapidly. When the 'A' value was about 200 pounds P_2O_5 per acre 50 percent of the soils showed benefit from phosphorus. With more than 300 pounds P_2O_5 per acre only 3 percent of the soils responded to phosphorus. Thus, the 'A' value will predict accurately the crop behavior when the 'A' values are either low or high, but the predic-

⁵This greenhouse experiment was conducted by Albert Hunter and Lawrence Alban, Oregon State College, Corvallis.




FIGURE 3.—Relationship for greenhouse and field experiments between percentage of soils showing a yield increase from phosphorus and 'A' value.

tion is uncertain when the 'A' value is around 200 pounds P₂O₅ per acre.

If the crop yield on the untreated soils is expressed as a percentage of the maximum yield, or yield from largest P₂O₅ rate used, a quantitative relationship with the 'A' value can be obtained by correlation analysis for the number of soils showing significant increases in crop yield from phosphate fertilization. The linear relationship between the 'A' value and percentage of maximum yield for each greenhouse experiment is shown in table 4. All the correlation coefficients are highly significant at the 1-percent level, but the r^2 values indicate that a large proportion of the variance in percentage of maximum yield is related to factors other than 'A' value.

A method for determining the phosphate-fertility level is most useful if it can be used to predict the crop behavior to additions of fertilizer. The standard error of estimate and the r^2 values (table 4) indicate that only a fair degree of precision may be expected for predicting percentage of maximum yield from 'A' values. The predictions would be expected to be more accurate for soils similar in the physical and chemical properties affecting plant growth and less accurate when these properties differ greatly. For example, the elimination of one soil from experiment 1 increases the r value from 0.690 to 0.721. The lack of high precision in predicting percentage of maximum yield from 'A' values does not necessarily mean the 'A' value fails to measure the phosphate-fertility level in the soil, because there may have been uncontrolled factors other than available phosphorus affecting plant growth.

TABLE 4.—*Linear correlations between 'A' values and percentage of maximum yield on the soils showing a significant increase in yield from added phosphorus*

Experiment No.	Number of soils	Correlation coefficient (<i>r</i>)	Coefficient of determination (<i>r</i> ²)×100	Standard error of estimate
1.....	43	0.690	48	21.4
2.....	34	.841	71	14.3
3.....	21	.758	57	13.8
4.....	22	.798	64	15.3
5.....	16	.828	69	9.5

The question arises as to whether the true relationship between percentage of maximum yield and the 'A' value is linear or curvilinear. The percentage figures were converted to angles (25, p. 449) and also used in the form of $\log(A-y)$, where A =maximum yield, or 100, and y =percentage of the maximum yield (3, 24). These transformations showed either a slight decrease, no change, or a slight increase in the correlation values with respect to the various experiments. Percentage-of-maximum-yield figures larger than 89 were not used, since the yield increase was not significantly different from the untreated soil. If percentage-yield figures between 89 and 98 are included, the logarithmic relationship, $\log(A-y)$, shows the best fit of the data plotted against 'A' values; however, the accuracy in measurement in this range is very poor. The linear relationship between percentage of maximum yield and 'A' values within the range of values indicated previously appears to fit the data equally as well as a curvilinear relationship.

From a practical standpoint, the prediction from 'A' value as to whether or not a yield response may be expected meets most of the requirements of a test to determine phosphate-fertility levels. The prediction from 'A' values of the percentage of maximum yield is a further refinement, which may be satisfactory on a limited group of soils; but, until the various factors affecting plant growth can be controlled better, any attempt to apply this relationship to a large number of soils will decrease the accuracy of the prediction. The maximum yield, for example, ranged from 10 to 31 gm. dry weight in experiment 1 under conditions where N, P₂O₅, and K₂O were added in amounts considered to be sufficient for maximum growth.

Field Experiments

Data on crop response to added phosphate in the field were available from 66 locations from experiment 1 and 29 samples from experiments 4 and 5. The relationship between 'A' value and the percentage of soils responding to phosphate is indicated by the broken line in figure 3. When the 'A' value was less than 100 pounds P₂O₅ per acre, 85 percent of the soils gave a phosphate response. None of the soils gave a phosphate response when the 'A' value was above 150 pounds P₂O₅ per acre, if a response for both field and greenhouse experiments were taken as a criterion of response. Actually, there

were 3 dryland wheat soils from Canada and 2 from Washington on which a field response but no response in the greenhouse was obtained. Four other soils from various locations also fell in this category.

Comparing soils from the field and greenhouse experiments, a response to phosphate was measurable in the greenhouse at higher levels of 'A' values than in the field (fig. 3). This result would be expected as a result of the smaller soil volume available to the plant roots in a greenhouse pot compared with field conditions.

MEASUREMENT OF NaHCO_3 -SOLUBLE PHOSPHORUS IN RELATION TO YIELD RESPONSE

All the soils responded to added phosphate in the greenhouse and 90 percent of them in the field when the extractable phosphorus was less than 25 pounds P_2O_5 per acre (fig. 4). When the extractable phosphorus was between 25 and 50 pounds P_2O_5 , 90 percent of the soils responded to phosphate in the greenhouse in contrast to only 50 percent in the field. These differences between field and greenhouse response are similar to those noted previously for the 'A' value comparisons in figure 3, and probably they are the result of the same effects.




FIGURE 4.—Relationship between NaHCO_3 -soluble phosphorus and the percentage of soils responding to phosphate in field and greenhouse experiments. The soils are the same as those compared in figure 3.

A similar comparison between field and greenhouse response was reported by Jenny and coworkers (15) for the relationship between the percentage of maximum yield of lettuce in pots and the number of field tests responding to phosphate. Most of the field tests responded to phosphate when the percentage of maximum yield of lettuce was less than 20 percent. When the percentage of maximum yield of lettuce was 40 percent or more, only a small percentage of the field tests responded to phosphate.

The average value of NaHCO_3 -soluble phosphorus for the 33 soils showing a phosphate response in the field (fig. 4) was 26 pounds P_2O_5 per acre. The values ranged from 15 to 53 pounds P_2O_5 per acre. The average 'A' value for these same soils was 80 pounds P_2O_5 per acre, with a range of 45 to 137.

CORRELATION OF 'A' VALUES WITH VARIOUS METHODS

The amount of phosphorus extracted by the various methods was correlated with 'A' values for six groups of soils (table 5), although not all of the methods were tested for each group. The NaHCO_3 method consistently gave a high correlation with 'A' value for all groups of soils, whereas the other methods were either lower than the NaHCO_3 method or variably correlated with the different groups of soils. The r^2 values for the 74 samples in experiment 1 were 0.882, 0.562, and 0.059 for the NaHCO_3 , Bray, and CO_2 methods, respectively. This comparison supplies convincing evidence of the close relationship between 'A' values and NaHCO_3 -soluble phosphorus for a group of soils containing both acid and calcareous types.

TABLE 5.—Correlations between the amount of phosphorus extracted by various methods and 'A' values in 6 experiments

Method	Correlation coefficient (r) for experiment—					
	1	2	3	4	5	6
NaHCO_3	0.939	0.903	0.874	0.936	0.935	0.866
Bray ¹750	.940	.802	.903	.898	-----
H_2O	-----	-----	.716	.965	.837	.751
CO_2243	-----	.628	.881	.774	.633

¹ Extracting solution is 0.025 N HCl plus 0.03 N NH_4F .

A further breakdown of the 74 soils based on their cation exchange-anion exchange (C/A) ratios is possible. According to Mehlich (20) the C/A is a qualitative measure for distinguishing between clay minerals of the 2:1 and 1:1 lattice types. When the C/A is greater than 2 or less than 1, the clay minerals are predominately 2:1 and 1:1, respectively. Previous experience with soil-test methods has led to the view that a given method is suitable only to soils with similar chemical and physical properties; however, when such separations into categories are made, the usual basis is the presence or absence of calcium carbonate.

The CO_2 method differs in its relationship to 'A' values as the C/A changes, but the NaHCO_3 , Bray, and Truog methods are not appreciably affected (table 6). The NaHCO_3 method shows the highest correlation with 'A' value for all three groups of soils separated according to C/A ratios.

The main objective of a soil test for available phosphorus is to estimate the phosphorus-fertility level. Assuming that the 'A' value




FIGURE 5.—Relationship between NaHCO_3 -soluble phosphorus and 'A' values.

is a measure of the phosphorus-fertility level in a soil, the accuracy with which 'A' values may be predicted from NaHCO_3 -soluble phosphorus can be determined from the regression line relating these two variables. The relationship between these 2 values for 212 samples (samples with 'A' values larger than 450 are not shown in the graph, but they are included in the regression equation) for the first 5 greenhouse experiments, is shown in figure 5. The correlation coefficient was 0.887 for the combined 212 samples. The equation of the regression line minimizing the Y components of deviation is $Y=2.05X+44$, and the standard error of the estimate is ± 69 pounds P_2O_5 per acre. The equation for the regression of NaHCO_3 values on 'A' values is $X=0.384Y+1.7$.

TABLE 6.—Effect of cation exchange-anion exchange ratios on the relationship between the amount of soluble phosphorus and 'A' values on 74 soils, experiment 1

Cation-anion exchange ratio	Number of soils	Correlation coefficient (r) for method ¹ —			
		CO_2	Truog	Bray	NaHCO_3
C/A, <1:1-----	16	0.096	0.661	0.772	0.917
C/A, 1:1 to 2:0-----	19	.297	.553	.798	.954
C/A, >2:1-----	39	.463	.694	.763	.953
	74	.243	.548	.750	.939

¹ Correlations for the CO_2 and Bray methods were supplied through the courtesy of J. W. Fitts; C/A ratios were determined by A. Mehlich.

More than 94 percent of the soils on which the plants responded to phosphate in the greenhouse have 'A' values less than 250 (fig. 3), so the major interest in 'A' values will be in this range. In addition, the error of measurement for 'A' values is larger for the high than for the low values. An error estimate for 'A' values from NaHCO_3 values based on a wide range of 'A' values would be too high for the low and too low for the high ranges of measurements. Therefore, the error of estimate for predicting 'A' values from NaHCO_3 soluble phosphorus was determined from the number of soils showing 'A' values less than 250 as well as for the total number of comparisons. This equation, $Y=1.7X+45$, is calculated from the 159 comparisons where the 'A' value is less than 250. The standard error of the estimate is ± 30 pounds P_2O_5 per acre.

DISCUSSION

A useful chemical extractant for plant available phosphorus ('A') in soils will (1) measure all or a definite proportion of the various forms of phosphorus in the same relative amounts as they are absorbed by plants during the growing season, (2) correlate to a high degree with plant uptake of phosphorus and yield response to added phosphorus over a wide range of soil types, (3) minimize the secondary

precipitation and adsorption reactions that may occur during the extraction, and (4) be adaptable to rapid, routine-test procedures.

Some of the currently used soil tests for phosphorus satisfy requirements (3) and (4) but fall short in varying degrees for (1) and (2). The chief incentive to develop a better test is the failure of a given method to apply to a wide variety of soils. If the limits of soil conditions and plant growth under which a given test works are known, satisfactory results may be expected, but too often these limitations are not adequately defined. The large number of soil-test methods used by various agencies not only indicates that each test will apply to a limited range of soils but also strongly implies a lack of agreement as to which test is best for a given group of soils with similar chemical and physical properties.

In this study attempts were made to find a chemical extractant that measured not only a definite proportion of the various forms of phosphorus available to plants but also minimized the secondary precipitation reactions occurring during the extraction process. The data presented indicate that the two major mechanisms of the extraction process are: (1) Calcium phosphates increase in solubility in NaHCO_3 at pH 8.5, as a result of the repression of the Ca^{++} activity (commonion effect of CO_3^{--} ions in the presence of solid phase CaCO_3); and (2) the HCO_3^- , CO_3^{--} , and OH^- ions replace phosphate ions on the surface of the soil particles. The phosphorus available to plants is considered to be derived chiefly from various forms of calcium phosphates and from the phosphorus on the surface of soil particles that readily exchanges with P^{32} in the soil solution. Probably a large proportion of the calcium phosphates available to plants is also present on the surface of the soil particles (21).

Acid extractants, such as CO_2 , weak organic acids, or mineral acids, are likely to be affected by secondary reactions on calcareous soils, whereas the NaHCO_3 method is essentially free from secondary reactions. To illustrate this difference, a calcareous soil-water system saturated with CO_2 will show a pH near 6.1 (27) if the CaCO_3 is in excess. When the soil-to-water ratio is 1:10 the CaCO_3 content must exceed 0.9 percent to be in excess, if the system is in equilibrium with 1 atmosphere of CO_2 . If the CaCO_3 content of the soil is less than 0.9 percent, all of the CaCO_3 will dissolve and the pH will decrease below 6.1. As a result, the extractions will not be made at the same pH, but will depend on the CaCO_3 content of the soil. The pH of the NaHCO_3 solution does not change when shaken with soil. These effects, in addition to those noted previously under the heading "Theoretical Basis," page 2, probably contribute to the lower correlation values found between CO_2 -soluble phosphorus and 'A' values.

Evidently the Bray method is less subject on calcareous soils to interfering secondary reactions than the CO_2 method. This difference is probably accounted for by the short extraction period used in the Bray procedure rather than by any essential differences in the chemical reactions involved. The extraction period is 5 minutes for the Bray procedure compared with 20 to 30 minutes for the CO_2 method. According to Burd (5) a short extraction period reduces secondary precipitation reactions of calcium phosphates in calcareous soils.

The lower correlation values (table 5) for the Bray method compared to NaHCO_3 on the calcareous soils are associated with a tendency for the Bray method to give low values for extractable phosphorus on soils high in CaCO_3 .

Water-soluble phosphate has been used in California with considerable success as an indicator of crop response (2, 19). In this study the water-soluble phosphate gave a higher correlation with 'A' values than NaHCO_3 in one experiment, but a considerably lower correlation than NaHCO_3 in three other experiments. The data indicate that the water-soluble phosphate tends to give lower correlations with 'A' values as the number of soil types compared increases.

RECOMMENDATIONS

The level of phosphorus soluble in NaHCO_3 , at which most of the soils respond to applied phosphate, is lower for the field than for the greenhouse experiments (fig. 4). Based on field experiments, using a 1:20 soil-to-solution ratio, the relationship between expected yield response to applied phosphate and the level of NaHCO_3 -soluble phosphorus is as follows: Below 25 pounds P_2O_5 per acre, a response; 25 to 50 pounds, a probable response; above 50 pounds, a response unlikely. These values were obtained from experiments using wheat, oats, and alfalfa. The values may be expected to apply also for crops with similar phosphorus requirements, but the values may need to be lower for corn and higher for potatoes.

SUMMARY

A new method for extracting soil phosphorus with 0.5 molar solutions of NaHCO_3 at pH 8.5 is described. The method is adaptable to rapid, routine, soil-testing procedures.

A theoretical basis for development of the method is described. The extraction mechanism is based on the increased solubility of calcium phosphates as a result of lowering the Ca^{++} activity in solution. Also, the NaHCO_3 extracts about one-half of the amount of phosphorus on the surface of soil particles, which readily exchanges with P^{32} in the soil solution.

Secondary precipitation and adsorption reactions are greatly minimized in the NaHCO_3 extractions.

Evaluation of the method is based on 212 soil samples from greenhouse and 95 soil samples from field experiments. The relationships between yield response of crops to phosphate and 'A' values determined in the greenhouse experiments are indicated.

The NaHCO_3 method was compared with the Bray, Truog, water-soluble, and CO_2 -soluble phosphorus methods by means of correlation analysis with 'A' values from six experiments. The NaHCO_3 method gave the highest correlations with 'A' values with two exceptions—one compared to the Bray and one compared to the water-soluble methods.

Recommendations are given concerning the relation of the expected crop response in the field to the level of phosphorus soluble in NaHCO_3 .

LITERATURE CITED

- (1) BENNE, E. J., PERKINS, A. T., and KING, H. H.
1936. THE EFFECT OF CALCIUM IONS AND REACTION UPON THE SOLUBILITY OF PHOSPHORUS. *Soil Sci.* 42: 29-38, illus.
- (2) BINGHAM, F. T.
1949. SOIL TEST FOR PHOSPHATE. . . . *Calif. Agr.* 3 (8): 11, 14, illus.
- (3) BRAY, R. H.
1948. CORRELATION OF SOIL TESTS WITH CROP RESPONSE TO ADDED FERTILIZERS AND WITH FERTILIZER REQUIREMENT. *In Diagnostic Techniques for Soils and Crops*, pp. 53-86, illus. American Potash Institute, Washington, D. C.
- (4) ——— and KURTZ, L. T.
1945. DETERMINATION OF TOTAL, ORGANIC, AND AVAILABLE FORMS OF PHOSPHORUS IN SOILS. *Soil Sci.* 59: 39-45.
- (5) BURD, J. S.
1948. CHEMISTRY OF THE PHOSPHATE ION IN SOIL SYSTEMS. *Soil Sci.* 65: 227-247, illus.
- (6) ——— and MURPHY, H. F.
1939. THE USE OF CHEMICAL DATA IN THE PROGNOSIS OF PHOSPHATE DEFICIENCY IN SOILS. *Hilgardia* 12: 323-340, illus.
- (7) COLE, C. V., OLSEN, S. R., and SCOTT, C. O.
1953. THE NATURE OF PHOSPHATE ADSORPTION BY CALCIUM CARBONATE. *Soil Sci. Soc. Amer. Proc.* 17: 352-356, illus.
- (8) COOKE, G. W.
1951. FIXATION OF PHOSPHATE DURING THE ACID EXTRACTION OF SOILS. *Soil Sci. Jour.* 2: 254-262.
- (9) DAS, S.
1930. AN IMPROVED METHOD FOR THE DETERMINATION OF AVAILABLE PHOSPHORIC ACID OF SOILS. *Soil Sci.* 30: 33-48.
- (10) DICKMAN, S. R., and BRAY, R. H.
1940. COLORIMETRIC DETERMINATION OF PHOSPHATE. *Indus and Engin: Chem., Analyt. Ed.* 12: 665-668.
- (11) FREAR, G. L., and JOHNSTON, J.
1929. THE SOLUBILITY OF CALCIUM CARBONATE (CALCITE) IN CERTAIN AQUEOUS SOLUTIONS AT 25° C. *Amer. Chem. Soc. Jour.* 51: 2082-2093, illus.
- (12) FRIED, M., and DEAN, L. A.
1952. A CONCEPT CONCERNING THE MEASUREMENT OF AVAILABLE SOIL NUTRIENTS. *Soil Sci.* 73: 263-271.
- (13) GARDNER, R., and KELLEY, O. J.
1940. RELATION OF pH TO PHOSPHATE SOLUBILITY IN COLORADO SOILS: *Soil Sci.* 50: 91-102, illus.
- (14) HOCKENSMITH, R. D., GARDNER, R., and GOODWIN, J.
1933. COMPARISON OF METHODS FOR ESTIMATING AVAILABLE PHOSPHORUS IN ALKALINE CALCAREOUS SOILS. *Colo. Expt. Sta. Tech. Bul.* 2, 24 pp.
- (15) JENNY, H., VLAMIS, J., and MARTIN, W. E.
1950. GREENHOUSE ASSAY OF FERTILITY OF CALIFORNIA SOILS. *Hilgardia* 20: 1-8, illus.
- (16) KURTZ, T., DETURK, E. E., and BRAY, R. H.
1946. PHOSPHATE ADSORPTION BY ILLINOIS SOILS. *Soil Sci.* 61: 111-124, illus.
- (17) LAATSCH, W.
1941. DIE FESTLEGUNG DER PHOSPHORSÄURE AN TONE. *Bodenk. u. Pflanzenernähr.* 23: 17-31.
- (18) McGEORGE, W. T., BUEHRER, T. F., and BREAZEALE, J. F.
1935. PHOSPHATE AVAILABILITY IN CALCAREOUS SOILS: A FUNCTION OF CARBON DIOXIDE AND pH. *Amer. Soc. Agron. Jour.* 27: 330-335, illus.
- (19) MARTIN, W. E., and BUCHANAN, J. R.
1950. PHOSPHATE TEST FOR GRAIN LAND. *Calif. Agr.* 4 (12): 7, 12, illus.
- (20) MEHLICH, A.
1948. DETERMINATION OF CATION- AND ANION-EXCHANGE PROPERTIES OF SOILS. *Soil Sci.* 66: 429-445, illus.

- (21) OLSEN, S. R.
1953. INORGANIC PHOSPHORUS IN ALKALINE AND CALCAREOUS SOILS. Amer. Soc. Agron. Jour. Monog. 4: 89-122.
- (22) PRATT, P. F., and THORNE, D. W.
1948. SOLUBILITY AND PHYSIOLOGICAL AVAILABILITY OF PHOSPHATE IN SODIUM AND CALCIUM SYSTEM. Soil Sci. Soc. Amer. Proc. 13: 213-217, illus.
- (23) RHOADES, H. F.
1939. EFFECT OF ORGANIC MATTER DECOMPOSITION ON THE SOLUBILITY AND FIXATION OF PHOSPHORUS IN ALKALINE SOILS. Nebr. Agr. Expt. Sta. Res. Bul. 113, 23 pp.
- (24) SMITH, F. W., and COOK, R. L.
1953. A STUDY OF THE RELATIONSHIP BETWEEN CHEMICALLY AVAILABLE PHOSPHORUS AND PLANT GROWTH RESPONSE ON SEVERAL MICHIGAN SOILS. Soil Sci. Soc. Amer. Proc. 17: 26-30, illus.
- (25) SNEDECOR, G. W.
1946. STATISTICAL METHODS. Ed. 4, 485 pp., illus. Ames, Iowa.
- (26) STANBERRY, C. O.
1948. THE BEHAVIOR OF PHOSPHORUS IN AN ALKALINE IRRIGATED SOIL IN WASHINGTON. Soil Sci. Soc. Amer. Proc. 13: 205-212, illus.
- (27) WHITNEY, R. S., and GARDNER, R.
1943. THE EFFECT OF CARBON DIOXIDE ON SOIL REACTION. Soil Sci. 55: 127-141, illus.

Hortigro, LTD

Pukekohe, New Zealand
Phone +64 923 71777

Client: ADRIAN BALLE

City : AUCKLAND, NEW ZEALAND

Date : 7-Dec-18


Location		EHENSON		Previous Analyses & Applications						
Crop		PASTURE / PASTURE		12/20/2017		12/15/2016		12/4/2015		
Field / Sample		LUCERNE		13.91		9.74		9.82		
Lab No.		D0016		5.9		6.0		5.8		
Total Exchange Capacity (M.E.)		14.75		8.4		11.9		6.4		
Desired Ca : Mg, Percent		68 : 12								
pH of Soil Sample		5.6								
Humus Content, Percent		8.6								
BASE SATURATION PERCENT				%		%		%		
Calcium (60 to 70%)		50.10		61.57		66.43		61.97		
Magnesium (10 to 20%) } 80%		12.32		10.48		9.37		7.81		
Potassium (2 to 5%)		3.46		2.97		1.79		1.71		
Sodium (.5 to 3%)		1.50		1.34		1.94		1.70		
Other Bases (Variable)		6.22		5.64		5.47		5.81		
EXCHANGEABLE HYDROGEN (10 to 15%)		26.40		18.00		15.00		21.00		
		RECOMMENDATIONS								
S O I L I O N S	NITROGEN		Amendment		kg/ha		kg/ha		kg/ha	
	kg/ha	ENR Value	132	APPLY NITROGEN AS NEEDED						
	SULFATE - S		Value Found		28		Val Found 18	Val Found 20	Val Found 24	
p.p.m.		39		SULFUR 90-92%						
PHOSPHATES		Desired Value		280		Val Found 31	Val Found 267	Val Found 212		
as (P2O5)		Olsen Value		DAP 18-46-0		Val Found 239	Val Found 267	Val Found 212		
kg/ha		Value Found								
Deficit/Surplus		-340								
C A T I O N S	CALCIUM		PARKER LIME #1222 (a)		2186		Amend	added	Amend	added
	kg/ha	Desired Value	4497							
		Value Found	3313							
		Deficit/Surplus	-1184							
MAGNESIUM		KIESERIT 16% Mg (b)		925		(Apply 1/3 per year for next 3 years if lime is applied as recommended.)				
kg/ha	Desired Value	476								
	Value Found	489								
	Deficit/Surplus	+13								
POTASSIUM		POTASSIUM SULFATE		224						
kg/ha	Desired Value	644								
	Value Found	446								
	Deficit/Surplus	-198								
SODIUM						P.P.M		P.P.M		P.P.M
kg/ha	Desired Value	76								
	Value Found	114								
	Deficit/Surplus	+38								
T R A C E S	Boron	p.p.m.	0.96	BORON 14.3%	17	0.89	0.61	0.49		
	Iron	p.p.m.	15.69	FE SULFATE 21% (c) (d) (e)	325	16	21	14		
	Manganese	p.p.m.	19.99	MANG SULF 28% (f)	168	29	29	33		
	Copper	p.p.m.	0.18	CU SULFATE 23% (g)	39	0.39	0.67	0.70		
	Zinc	p.p.m.	4.25	ZINC SULFATE 36%	22	3.46	3.39	3.60		
							0.40	0.10	0.05	
N O T E S	(a) Apply Limestone at this rate only if no Lime or Gypsum has been added in the last three years, if previously limed we need to revise rates accordingly.									
	(b) Apply 1/3 of this amount of Kieserit 16% Mg each year for 3 years ONLY if the full amount of recommended lime is applied.									
	(c) Ferrous Sulfate - should only use the white or blue-green in color, black or rust colored is never recommended. CAUTION: Apply so as to avoid getting the dust from the material on the leaves of actively growing plants									
	(d) CAUTION: Only apply at this rate if yellowing or stunting is evident now or in past crops, otherwise test subsoil for Iron levels or use on a small test plot to establish need.									
	(e) WARNING: Apply at least one good rain or irrigation before liming is done.									
	(f) Apply this amount of Manganese Sulfate ONLY if recommended Ferrous Sulfate is applied, or if subsoil tests verify adequate iron is present there.									
	(g) Or can apply Copper Sulfate 23% at the rate of 5.6 kg/ha per year for 7 years.									

Target soil test ranges

Average calibration curves (bold centre line) are presented for each major soil group. The thinner line beside the solid average curve indicates that there is a 95% probability that the average relationship lies within this band. This applies to all the calibration curves. Target soil test levels are presented as a range for each major soil group.


Soil Olsen P

Sedimentary soil



The relationship between relative pasture production and Olsen P for **sedimentary** soils. The shaded box represents the target range for soil Olsen P.

Ash soils



The relationship between relative pasture production and Olsen P test for **ash** soils. The shaded box represents the target range for soil Olsen P.

Acknowledgments

The results in this booklet are based on comprehensive soil fertility and fertiliser research. The work of field researchers, past and present, who have conducted field trials under the auspices of the Agricultural Research Division (MAF), MAF Technology and latterly AgResearch is gratefully recognised.

The New Zealand Society of Animal Production gave permission for the use of its data on critical mineral concentrations.

The fertiliser manufacturing companies (Ballance Agri-Nutrients Limited and Ravensdown Fertiliser Co-operative Limited) have provided the funding for this revised edition. Additional copies are available from Ballance and Ravensdown



This booklet is published by the New Zealand Fertiliser Manufacturers' Research Association. For more information:

PO Box 9577, Newmarket, Auckland, New Zealand

Phone: 09 476 3079 Fax: 09 476 3059

Email: info@fertresearch.org.nz

Website: www.fertresearch.org.nz

First Produced by New Zealand Fertiliser Manufacturers' Research Association Inc. and New Zealand Pastoral Agriculture Research Institute Ltd, 1994

Revised 1999

Reprinted 2004

Revised 2009

©Notice of Copyright

All rights reserved. No part of this publication may be reproduced or transmitted, in any form by any means, electronic, mechanical, photocopying, recording or otherwise, or stored in any retrieval system of any nature without the prior written permission of the copyright holders and the publisher, application for which shall be made to the publisher/copyright holder. New Zealand Fertiliser Manufacturers' Research Association Inc. reserves the right to make changes to the information in this publication without notice.

ISBN 978-0-9864528-1-9